



PESANDOR

Variety obtained by selective breeding of the Jersey shallot, OYDV-virus free

Name of breeder: INRA
Name of maintainer: INRA
Botanical name: *Allium cepa* gr. aggregatum
Common name: Shallot
Variety name: PESANDOR
Variety type: Long

VARIETY INFORMATION SHEET



OBSERVED CHARACTERISTICS

FOLIAGE

| | | |
|---------------------------|----------------|---|
| Attitude | Semi-erect | 3 |
| Waxiness | Weak to medium | 4 |
| Intensity of green colour | Medium to dark | 6 |
| Cranking | Absent or weak | 1 |
| Length | Medium to long | 6 |
| Diameter | Medium | 5 |

BULB

| | | |
|--|---------------------|---|
| Number of bulblets (degree of splitting into bulblets with dry skin around each bulblet) | Medium to strong | 6 |
| Bulblet size | Small | 3 |
| Bulblet height | Tall to very tall | 8 |
| Bulblet diameter | Very small to small | 2 |
| Position of maximum diameter | In middle | 2 |
| Width of neck | Narrow | 3 |
| Shape in longitudinal section | Medium ovate | 2 |
| Shape of stem end | Strongly sloping | 6 |
| Shape of root end | Strongly tapered | 5 |
| Base colour of dry skin | Brown | 5 |
| Intensity of base colour of dry skin (intensity of colour) | Medium to dark | 6 |
| Secondary colour of skin (= hue of dry skin) | Brownish | 5 |
| Colour of epidermis of fleshy scales | Reddish | 3 |
| Colour of flesh | Pinkish | 3 |
| Intensity of flesh colour | Dark | 7 |

PHYSIOLOGICAL CHARACTERISTICS

| | | |
|--|---------------------|---|
| Number of growing points per kg | Medium | 5 |
| Dry matter content | Medium to high | 6 |
| Tendency to bolt | Absent or very weak | 1 |
| Time of harvest maturity | Early | 3 |
| Time of sprouting during storage (=storage period) | Late | 7 |
| Male sterility | Absent or very weak | 1 |